



Cervical Cancer

What is cervical cancer?


Cancer starts when cells in the body begin to grow out of control. Cells in nearly any part of the body can become cancer, and can spread to other areas of the body. To learn more about how cancers start and spread, see [What Is Cancer?](#)

Cervical cancer starts in the cells lining the cervix -- the lower part of the uterus (womb). This is sometimes called the uterine cervix. The fetus grows in the body of the uterus (the upper part). The cervix connects the body of the uterus to the vagina (birth canal).

The cervix has two different parts and is covered with two different types of cells.

- The part of the cervix closest to the body of the uterus is called the **endocervix** and is covered with **glandular** cells.
- The part next to the vagina is the **exocervix** (*or* **ectocervix**) and is covered in **squamous** cells.

These two cell types meet at a place called the **transformation zone**. The exact location of the transformation zone changes as you get older and if you give birth.



Most cervical cancers begin in the cells in the transformation zone. These cells do not suddenly change into cancer. Instead, the normal cells of the cervix first gradually develop pre-cancerous changes that turn into cancer. Doctors use several terms to describe these pre-cancerous changes, including *cervical intraepithelial neoplasia (CIN)*, *squamous intraepithelial lesion (SIL)*, and *dysplasia*. These changes can be detected by the Pap test and treated to prevent cancer from developing (see [Can cervical cancer be prevented?](#)).

Although cervical cancers start from cells with pre-cancerous changes (pre-cancers), only some of the women with pre-cancers of the cervix will develop cancer. It usually takes several years for cervical pre-cancer to change to cervical cancer, but it also can happen in less than a year. For most women, pre-cancerous cells will go away without any treatment. Still, in some women pre-cancers turn into true (invasive) cancers. Treating all cervical pre-cancers can prevent almost all cervical cancers. Pre-cancerous changes and specific types of treatment for pre-cancers are discussed in [Cervical Cancer Prevention and Early Detection](#).

Types of cervical cancer

Cervical cancers and cervical pre-cancers are classified by how they look under a microscope. The main types of cervical cancers are *squamous cell carcinoma* and *adenocarcinoma*.

- Most (up to 9 out of 10) cervical cancers are **squamous cell carcinomas**. These cancers develop from cells in the exocervix and the cancer cells have features of squamous cells

under the microscope. Squamous cell carcinomas most often begin in the transformation zone (where the exocervix joins the endocervix).

- Most of the other cervical cancers are **adenocarcinomas**. Adenocarcinomas are cancers that develop from gland cells. Cervical adenocarcinoma develops from the mucus-producing gland cells of the endocervix. Cervical adenocarcinomas seem to have become more common in the past 20 to 30 years.
- Less commonly, cervical cancers have features of both squamous cell carcinomas and adenocarcinomas. These are called **adenosquamous carcinomas** or **mixed carcinomas**.

Although almost all cervical cancers are either squamous cell carcinomas or adenocarcinomas, other types of cancer also can develop in the cervix. These other types, such as melanoma, sarcoma, and lymphoma, occur more commonly in other parts of the body.

Only the more common cervical cancer types are covered here, and not the rare types.

What are the key statistics about cervical cancer?

The American Cancer Society's estimates for cervical cancer in the United States for 2017 are:

- About 12,820 new cases of invasive cervical cancer will be diagnosed.
- About 4,210 women will die from cervical cancer.

Cervical pre-cancers are diagnosed far more often than invasive cervical cancer.

Cervical cancer was once one of the most common causes of cancer death for American women. But over the last 40 years, the cervical cancer death rate has gone down by more than 50%. The main reason for this change was the increased use of the Pap test. This screening procedure can find changes in the cervix before cancer develops. It can also find cervical cancer early – when it is in its most curable stage.

Most cases of cervical cancer are found in women younger than 50. It rarely develops in women younger than 20. Many older women do not realize that the risk of developing cervical cancer is still present as they age. About 20% of cervical cancers are found in women older than 65. These cancers, however, rarely occur in women who have been getting regular tests for cervical cancer before they were 65. See [Can cervical cancer be prevented?](#) and [Cervical Cancer Prevention and Early Detection](#) for more information about tests used to screen for cervical cancer.

In the United States, Hispanic women are most likely to get cervical cancer, followed by African-Americans, American Indians and Alaskan natives, and whites. Asians and Pacific Islanders have the lowest risk of cervical cancer in this country.

Visit the American Cancer Society's Cancer Statistics Center for more key statistics.

What are the risk factors for cervical cancer?

A risk factor is anything that changes your chance of getting a disease such as cancer. Different cancers have different risk factors. For example, exposing skin to strong sunlight is a risk factor for skin cancer. Smoking is a risk factor for many cancers. But having a risk factor, or even several, does not mean that you will get the disease.

Several risk factors increase your chance of developing cervical cancer. Women without any of these risk factors rarely develop cervical cancer. Although these risk factors increase the odds of developing cervical cancer, many women with these risks do not develop this disease. When a woman develops cervical cancer or pre-cancerous changes, it might not be possible to say that a particular risk factor was the cause.

In thinking about risk factors, it helps to focus on those you can change or avoid (like smoking or human papilloma virus infection), rather than those you cannot (such as your age and family history). However, it is still important to know about risk factors that cannot be changed, because it's even more important for women who have these factors to get regular Pap tests to detect cervical cancer early.

Cervical cancer risk factors include:

Human papilloma virus (HPV) infection

Infection by the human papilloma virus (HPV) is the most important risk factor for cervical cancer. HPV is a group of more than 150 related viruses. Some of them cause a type of growth called **papillomas**, which are more commonly known as warts.

- HPV can infect cells on the surface of the skin, and those lining the genitals, anus, mouth and throat, but not the blood or internal organs such as the heart or lungs.
- HPV can spread from one person to another during skin-to-skin contact. One way HPV spreads is through sex, including vaginal, anal, and even oral sex.
- Different types of HPV cause warts on different parts of the body. Some cause common warts on the hands and feet; others tend to cause warts on the lips or tongue.

Certain types of HPV may cause warts on or around the female and male genital organs and in the anal area. These are called *low-risk types* of HPV because they are seldom linked to cancer.

Other types of HPV are called high-risk types because they are strongly linked to cancers, including cancer of the cervix, vulva, and vagina in women, penile cancer in men, and cancers of the anus, mouth, and throat in both men and women.

Doctors believe that a woman must be infected with HPV in order to develop cervical cancer. Although this can mean infection with any of the high-risk types, about two-thirds of all cervical cancers are caused by HPV 16 and 18.

Infection with HPV is common, and in most people the body can clear the infection by itself. Sometimes, however, the infection does not go away and becomes chronic. Chronic infection, especially when it is caused by certain high-risk HPV types, can eventually cause certain cancers, such as cervical cancer.

Although there is currently no cure for HPV infection, there are ways to treat the warts and abnormal cell growth that HPV causes.

For more information on about this topic, see [Cervical Cancer Prevention and Early Detection and HPV and HPV Testing](#).

Smoking

When someone smokes, they and those around them are exposed to many cancer-causing chemicals that affect organs other than the lungs. These harmful substances are absorbed through the lungs and carried in the bloodstream throughout the body.

Women who smoke are about twice as likely as non-smokers to get cervical cancer. Tobacco by-products have been found in the cervical mucus of women who smoke. Researchers believe that these substances damage the DNA of cervix cells and may contribute to the development of cervical cancer. Smoking also makes the immune system less effective in fighting HPV infections.

Immunosuppression

Human immunodeficiency virus (HIV), the virus that causes AIDS, damages a woman's immune system and puts them at higher risk for HPV infections. This might explain why women with AIDS have a higher risk for cervical cancer. The immune system is important in destroying cancer cells and slowing their growth and spread. In women with HIV, a cervical pre-cancer might develop into an invasive cancer faster than it normally would. Another group of women at risk for cervical cancer are those taking drugs to suppress their immune response, such as those being treated for an autoimmune disease (in which the immune system sees the body's own tissues as foreign and attacks them, as it would a germ) or those who have had an organ transplant.

Chlamydia infection

Chlamydia is a relatively common kind of bacteria that can infect the reproductive system. It is spread by sexual contact. Chlamydia infection can cause pelvic inflammation, leading to infertility. Some studies have seen a higher risk of cervical cancer in women whose blood tests and cervical mucus showed evidence of past or current chlamydia infection. Women who are infected with chlamydia often have no symptoms. In fact, they may not know that they are infected at all unless they are tested for chlamydia during a pelvic exam.

A diet low in fruits and vegetables

Women whose diets don't include enough fruits and vegetables may be at increased risk for cervical cancer.

Being overweight

Overweight women are more likely to develop adenocarcinoma of the cervix.

Long-term use of oral contraceptives (birth control pills)

There is evidence that taking oral contraceptives (OCs) for a long time increases the risk of cancer of the cervix. Research suggests that the risk of cervical cancer goes up the longer a woman takes OCs, but the risk goes back down again after the OCs are stopped, and returns to normal about 10 years after stopping.

The American Cancer Society believes that a woman and her doctor should discuss whether the benefits of using OCs outweigh the potential risks.

Intrauterine device use

Some research suggests that women who had ever used an intrauterine device (IUD) had a lower risk of cervical cancer. The effect on risk was seen even in women who had an IUD for less than a year, and the protective effect remained after the IUDs were removed.

Using an IUD might also lower the risk of endometrial (uterine) cancer. However, IUDs do have some risks. A woman interested in using an IUD should first discuss the possible risks and benefits with her doctor. Also, a woman with multiple sexual partners should use condoms to lower her risk of sexually transmitted illnesses no matter what other form of contraception she uses.

Having multiple full-term pregnancies

Women who have had 3 or more full-term pregnancies have an increased risk of developing cervical cancer. No one really knows why this is true. One theory is that these women had to have had unprotected intercourse to get pregnant, so they may have had more exposure to HPV. Also, studies have pointed to hormonal changes during pregnancy as possibly making women more susceptible to HPV infection or cancer growth. Another thought is that pregnant women might have weaker immune systems, allowing for HPV infection and cancer growth.

Being younger than 17 at your first full-term pregnancy

Women who were younger than 17 years when they had their first full-term pregnancy are almost 2 times more likely to get cervical cancer later in life than women who waited to get pregnant until they were 25 years or older.

Economic status

Many low-income women do not have easy access to adequate health care services, including Pap tests. This means they may not get screened or treated for cervical pre-cancers.

Diethylstilbestrol (DES)

DES is a hormonal drug that was given to some women between 1940 and 1971 to prevent miscarriage. Women whose mothers took DES (when pregnant with them) develop clear-cell adenocarcinoma of the vagina or cervix more often than would normally be expected. These types of cancer are extremely rare in women who haven't been exposed to DES. There is about 1 case of vaginal or cervical clear-cell adenocarcinoma in every 1,000 women whose mothers took DES during pregnancy. This means that about 99.9% of "DES daughters" do not develop these cancers.

DES-related clear cell adenocarcinoma is more common in the vagina than the cervix. The risk appears to be greatest in women whose mothers took the drug during their first 16 weeks of pregnancy. The average age of women diagnosed with DES-related clear-cell adenocarcinoma is 19 years. Since the use of DES during pregnancy was stopped by the FDA in 1971, even the youngest DES daughters are older than 40 – past the age of highest risk. Still, there is no age cut-off when these women are felt to be safe from DES-related cancer. Doctors do not know exactly how long these women will remain at risk.

DES daughters may also be at increased risk of developing squamous cell cancers and pre-cancers of the cervix linked to HPV.

You can learn more about DES in [DES Exposure: Questions and Answers](#). Read it on our website, or call (1-800-227-2345) to have a free copy sent to you.

Having a family history of cervical cancer

Cervical cancer may run in some families. If your mother or sister had cervical cancer, your chances of developing the disease are 2 to 3 times higher than if no one in the family had it. Some researchers suspect that some instances of this familial tendency are caused by an inherited condition that makes some women less able to fight off HPV infection than others. In other instances, women in the same family as a patient already diagnosed could be more likely to have one or more of the other non-genetic risk factors previously described in this section.

Do we know what causes cervical cancer?

In recent years, there has been a lot of progress in understanding what happens in cells of the cervix when cancer develops. In addition, several risk factors have been identified that increase the odds that a woman might develop cervical cancer (see [What Are the Risk Factors for Cervical Cancer](#)).

The development of normal human cells mostly depends on the information contained in the cells' DNA. DNA is the chemical in our cells that makes up our genes, which control how our cells work. We look like our parents because they are the source of our DNA. But DNA affects more than just how we look.

Some genes control when cells grow, divide, and die:

- Genes that help cells grow, divide, and stay alive are called **oncogenes**.
- Genes that help keep cell growth under control or make cells die at the right time are called **tumor suppressor genes**.

Cancers can be caused by DNA mutations (gene defects) that turn on oncogenes or turn off tumor suppressor genes.

HPV causes the production of two proteins known as E6 and E7 which turn off some tumor suppressor genes. This may allow the cervical lining cells to grow too much and to develop changes in additional genes, which in some cases will lead to cancer.

But HPV is not the only cause of cervical cancer. Most women with HPV don't get cervical cancer, and certain other risk factors, like smoking and HIV infection, influence which women exposed to HPV are more likely to develop cervical cancer.

Can cervical cancer be prevented?

The most common form of cervical cancer starts with pre-cancerous changes and there are ways to stop this disease from developing. The first way is to find and treat pre-cancers before they become true cancers, and the second is to prevent the pre-cancers.

Finding cervical pre-cancers

A well-proven way to prevent cervical cancer is to have testing (screening) to find pre-cancers before they can turn into invasive cancer. The Pap test (or Pap smear) and the human papilloma virus (HPV) test are used for this. If a pre-cancer is found it can be treated, stopping cervical cancer before it really starts. Most invasive cervical cancers are found in women who have not had regular Pap tests.

The Pap test is a procedure used to collect cells from the cervix so that they can be looked at under a microscope to find cancer and pre-cancer. These cells can also be used for HPV testing. A Pap test can be done during a pelvic exam, but not all pelvic exams include a Pap test.

An HPV test can be done on the same sample of cells collected from the Pap test.

The most important thing you can do to prevent cervical cancer is to be tested according to American Cancer Society guidelines. These can be found in Cervical Cancer Prevention and Early Detection. Information on treatment if the Pap test results are abnormal is also covered.

Things to do to prevent pre-cancers

There are also some things you can do to prevent pre-cancers, such as:

- Avoiding exposure to HPV
- Getting an HPV vaccine
- Not smoking

More information about ways to prevent cervical pre-cancer and cancer can be found in Cervical Cancer Prevention and Early Detection.

You can also find information on preventing HPV infection in HPV Vaccines.

Can cervical cancer be found early?

The best way to find cervical cancer early is to have regular screening with a Pap test (which may be combined with a test for human papilloma virus or HPV). As Pap testing became routine in this country, finding pre-invasive lesions (pre-cancers) of the cervix became far more common than finding invasive cancer. Being alert to any signs and symptoms of cervical cancer can also help avoid unnecessary delays in diagnosis. Early detection greatly improves the chances of successful treatment and can prevent any early cervical cell changes from becoming cancerous.

More information about using the Pap test and the HPV test to find cervical cancer early, including the American Cancer Society's Guidelines for cervical cancer screening can be found in Cervical Cancer Prevention and Early Detection.

Signs and symptoms of cervical cancer

Women with early cervical cancers and pre-cancers usually have no symptoms. Symptoms often do not begin until the cancer becomes invasive and grows into nearby tissue. When this happens, the most common symptoms are:

- Abnormal vaginal bleeding, such as bleeding after vaginal sex, bleeding after menopause, bleeding and spotting between periods, and having (menstrual) periods that are longer or heavier than usual. Bleeding after douching or after a pelvic exam may also occur.
- An unusual discharge from the vagina. The discharge may contain some blood and may occur between your periods or after menopause.
- Pain during sex.

These signs and symptoms can also be caused by conditions other than cervical cancer. For example, an infection can cause pain or bleeding. Still, if you have any of these symptoms, see a health care professional right away. Ignoring symptoms may allow the cancer to grow to a more advanced stage and lower your chance for effective treatment.

Even better, don't wait for symptoms to appear. Have regular Pap tests and pelvic exams.

Tests for cervical cancer

The first step in finding cervical cancer is often an abnormal Pap test result. This will lead to further tests which can diagnose cervical cancer.

Cervical cancer may also be suspected if you have symptoms like abnormal vaginal bleeding or pain during sex. Your primary doctor or gynecologist often can do the tests needed to diagnose pre-cancers and cancers and may also be able to treat a pre-cancer.

If there is a diagnosis of invasive cancer, your doctor should refer you to a gynecologic oncologist, a doctor who specializes in cancers of women's reproductive systems.

Tests for women with symptoms of cervical cancer or abnormal Pap results

Medical history and physical exam

First, the doctor will ask you about your personal and family medical history. This includes information related to risk factors and symptoms of cervical cancer. A complete physical exam will help evaluate your general state of health. The doctor will do a pelvic exam and may do a Pap test if one has not already been done. In addition, your lymph nodes will be felt for evidence of metastasis (cancer spread).

The Pap test is a screening test, not a diagnostic test. It cannot tell for certain if you have cervical cancer. An abnormal Pap test result may mean more testing, sometimes including tests to see if a cancer or a pre-cancer is actually present. The tests that are used include colposcopy (with biopsy), endocervical scraping, and cone biopsies.

Colposcopy

If you have certain symptoms that are suggestive of cancer or if your Pap test result shows abnormal cells, you will need to have a test called *colposcopy*. You will lie on the exam table as you do with a pelvic exam. A speculum will be placed in the vagina to help the doctor see the cervix. The doctor will use a colposcope to examine the cervix. The colposcope is an instrument that stays outside the body and has magnifying lenses. It lets the doctor see the surface of the cervix closely and clearly. Colposcopy itself is usually no more uncomfortable than any other speculum exam. It can be done safely even if you are pregnant. Like the Pap test, it is better not to do it during your menstrual period.

The doctor will put a weak solution of acetic acid (similar to vinegar) on your cervix to make any abnormal areas easier to see. If an abnormal area is seen, a biopsy (removal of a small piece of tissue) will be done. The tissue is sent to a lab to be looked at under a microscope. A biopsy is the best way to tell for certain if an abnormal area is a pre-cancer, a true cancer, or neither. Although the colposcopy procedure is usually not painful, the cervical biopsy can cause discomfort, cramping, bleeding, or even pain in some women.

Cervical biopsies

Several types of biopsies can be used to diagnose cervical pre-cancers and cancers. After these procedures, patients could feel mild cramping or pain and might also have some light bleeding.

Colposcopic biopsy

For this type of biopsy, the cervix is examined with a colposcope to find the abnormal areas. The cervix might be numbed with a local anesthetic before the biopsy. Using biopsy forceps, a small section of the abnormal area is removed.

Endocervical curettage (endocervical scraping)

Sometimes the transformation zone (the area at risk for HPV infection and pre-cancer) cannot be seen with the colposcope so something else must be done to check that area for cancer. This means inserting a narrow instrument (called a **curette**) into the endocervical canal (the part of the cervix closest to the uterus). The curette is used to scrape the inside of the canal to remove some of the tissue, which is then sent to the lab for examination.

Cone biopsy

In this procedure, also known as **conization**, the doctor removes a cone-shaped piece of tissue from the cervix. The tissue removed in the cone includes the transformation zone where cervical pre-cancers and cancers are most likely to start.

A cone biopsy is not only used to diagnose pre-cancers and cancers. It can also be used as a treatment since it can sometimes completely remove pre-cancers and some very early cancers.

The methods commonly used for cone biopsies are the loop electrosurgical excision procedure (LEEP), also called the **large loop excision of the transformation zone** (LLETZ), and the cold knife cone biopsy. With both procedures, you might have mild cramping and some bleeding for a few weeks.

- **Loop electrosurgical procedure (LEEP, LLETZ):** In this method, the tissue is removed with a thin wire loop that is heated by electricity and acts as a small knife. For this procedure, a local anesthetic is used, and it can be done in your doctor's office.
- **Cold knife cone biopsy:** This method is done in a hospital. A surgical scalpel or a laser is used to remove the tissue instead of a heated wire. You will receive anesthesia during the operation (either a general anesthesia, where you are asleep, or a spinal or epidural anesthesia, where an injection into the area around the spinal cord makes you numb below the waist). Having any type of cone biopsy will not prevent most women from getting pregnant, but if a large amount of tissue has been removed, women may have a higher risk of giving birth prematurely.

How biopsy results are reported

Pre-cancerous changes in a biopsy are called cervical intraepithelial neoplasia (CIN). Sometimes the term **dysplasia** is used instead of CIN. CIN is graded on a scale of 1 to 3 based on how much of the cervical tissue looks abnormal when viewed under the microscope.

- In CIN1, not much of the tissue looks abnormal, and it is considered the least serious cervical pre-cancer (mild dysplasia).
- In CIN2 more of the tissue looks abnormal (moderate dysplasia)
- In CIN3 most of the tissue looks abnormal; CIN3 is the most serious pre-cancer (severe dysplasia) and includes carcinoma in situ.

If a biopsy shows a pre-cancer, doctors will take steps to keep an actual cancer from developing. Treatment of women with abnormal pap results is discussed in Cervical Cancer Prevention and Early Detection.

If a cancer is found on a biopsy, it will be identified as either squamous cell carcinoma, adenocarcinoma, or another type.

Tests for women diagnosed with cervical cancer

If biopsy results show that you have cervical cancer, your doctor may order certain tests to see how far the cancer has spread. Many of the tests described below are not needed for every patient. Decisions about using these tests are based on the results of the physical exam and biopsy.

Cystoscopy and proctoscopy

These are most often done in women who have large tumors. They are often not needed if the cancer is caught early.

In **cystoscopy**, a slender tube with a light is placed into the bladder through the urethra. This lets the doctor check your bladder and urethra to see if cancer is growing into these areas. Suspicious areas that look like cancer can be biopsied during the procedure for testing. Cystoscopy can be done under a local anesthetic, but some women may need general anesthesia (where you are in a deep sleep). Your doctor will let you know what to expect before and after the procedure.

Proctoscopy is a procedure that lets the doctor see of the inside the rectum through a lighted tube to check for spread of cervical cancer into that area.

Your doctor may also do a pelvic exam while you are under anesthesia to find out if the cancer has spread beyond the cervix.

Imaging tests

If your doctor finds that you have cervical cancer, certain imaging tests may be done to look inside the body. These tests can show if and where the cancer has spread, which will help you and your doctor decide on a treatment plan.

Chest x-ray

Your chest may be x-rayed to see if cancer has spread to your lungs. This is very unlikely unless the cancer is far advanced.

Computed tomography (CT)

CT scans are usually done if the tumor is larger or if there is concern about cancer spread. For more information, see [CT Scan for Cancer](#).

Magnetic resonance imaging (MRI)

MRI looks at soft tissue parts of the body sometimes better than other imaging tests. Your doctor will decide which imaging test is best for your situation.

For more information, see [MRI for Cancer](#).

Intravenous urography

Intravenous urography (also known as **intravenous pyelogram**, or **IVP**) is an x-ray of the urinary system taken after a special dye is injected into a vein. This test can find abnormal areas in the urinary tract, caused by the spread of cervical cancer. The most common finding is a blockage of the ureters (tubes that connect the kidneys to the bladder) by the cancer. IVP is rarely used for patients with cervical cancer because CT and MRI are also good at finding abnormal areas in the urinary tract, as well as others not seen with an IVP.

Positron emission tomography (PET scan)

PET scans use glucose (a form of sugar) that contains a radioactive atom. Cancer cells in the body absorb large amounts of the radioactive sugar and a special camera can detect the radioactivity.

This test can help see if the cancer has spread to lymph nodes. PET scans can also be useful if your doctor thinks the cancer has spread but doesn't know where, because they scan your whole body.

PET scans are often combined with CT scans using a machine that can do both at the same time. The combined PET/CT test is rarely used for patients with early cervical cancer, but may be used to look for more advanced cancer or if radiation treatment is a possibility. For more information on this test, see [Nuclear Scans for Cancer](#).

Cervical cancer stages

A cancer's stage describes how far it has spread when it is first diagnosed. It is one of the key factors used by the patient's cancer team (of health care professionals) in determining the best treatment options. Information from exams and tests is used to determine the size of the

tumor, how deeply the tumor has invaded tissues in and around the cervix, and its spread to distant places (metastasis). For more information see Cancer Staging.

The **FIGO (International Federation of Gynecology and Obstetrics) staging system** is used most often for cancers of the female reproductive organs, including cervical cancer. Staging is based mainly on the results of the doctor's physical exam and a few other tests that are done in some cases, such as cystoscopy and proctoscopy. It is not based on what is found during surgery.

When surgery is done, it might show that the cancer has spread more than the doctors first thought. This new information could change the treatment plan, but it does not change the patient's stage.

The **AJCC (American Joint Committee on Cancer) TNM staging system** classifies cancer on the basis of 3 factors:

- The extent of the main **tumor (T)**
- Whether the cancer has spread to nearby lymph **nodes (N)**
- Whether the cancer has spread (**metastasized**) to distant parts of the body (**M**)

The FIGO system uses the same basic information as AJCC, except it does not use the lymph node status and does not include Stage 0 (carcinoma in situ) cervical cancer. Even though there are minor differences between these systems, the FIGO and TNM stages still closely match, and it is acceptable to use either one.

Tumor extent (T)

Tis: The cancer cells are only found on the surface of the cervix, without growing into deeper tissues. (Tis is not included in the FIGO system)

T1: The cancer cells have grown from the surface of the cervix into deeper tissues of the cervix. The cancer may also be growing into the body of the uterus, but it has not grown outside the uterus.

T1a: There is a very small amount of cancer, and it can be seen only under a microscope.

- **T1a1:** The area of cancer is less than 3 mm (about 1/8-inch) deep and less than 7 mm (about 1/4-inch) wide.
- **T1a2:** The area of cancer invasion is between 3 mm and 5 mm (about 1/5-inch) deep and less than 7 mm (about 1/4-inch) wide.

T1b: This stage includes stage I cancers that can be seen without a microscope. This stage also includes cancers that can only be seen with a microscope if they have spread deeper than 5 mm (about 1/5 inch) into connective tissue of the cervix or are wider than 7 mm.

- **T1b1:** The cancer can be seen but it is not larger than 4 cm (about 1 3/5 inches).
- **T1b2:** The cancer can be seen and is larger than 4 cm.

T2: In this stage, the cancer has grown beyond the cervix and uterus, but hasn't spread to the walls of the pelvis or the lower part of the vagina. The cancer may have grown into the upper part of the vagina.

T2a: The cancer has not spread into the tissues next to the cervix (called the **parametria**).

- **T2a1:** The cancer can be seen but it is not larger than 4 cm (about 1 3/5 inches).
- **T2a2:** The cancer can be seen and is larger than 4 cm.

T2b: The cancer has spread into the tissues next to the cervix (the parametria)

T3: The cancer has spread to the lower part of the vagina or the walls of the pelvis. The cancer may be blocking the ureters (tubes that carry urine from the kidneys to the bladder).

- **T3a:** The cancer has spread to the lower third of the vagina but not to the walls of the pelvis.
- **T3b:** The cancer has grown into the walls of the pelvis and/or is blocking one or both ureters causing kidney problems (called hydronephrosis).

T4: The cancer has spread to the bladder or rectum or it is growing out of the pelvis

Lymph node spread (N)

NX: The nearby lymph nodes cannot be assessed

N0: No spread to nearby lymph nodes

N1: The cancer has spread to nearby lymph nodes

Distant spread (M)

M0: The cancer has not spread to distant lymph nodes, organs, or tissues

M1: The cancer has spread to distant organs (such as the lungs or liver), to lymph nodes in the chest or neck, and/or to the peritoneum (the tissue coating the inside of the abdomen).

Stage grouping and FIGO stages

Information about the tumor (T), lymph nodes (N), and any cancer spread (M) is then combined to assign the cancer an overall stage. This process is called **stage grouping**. The stages are described using the number 0 and Roman numerals from I to IV. Some stages are divided into sub-stages indicated by letters and numbers. FIGO stages are the same as AJCC stages, except that stage 0 doesn't exist in the FIGO system.

Stage 0 (Tis, N0, M0)

The cancer cells are only in the cells on the surface of the cervix (the layer of cells lining the cervix), without growing into (invading) deeper tissues of the cervix. This stage is also called *carcinoma in situ* (CIS) which is part of cervical intraepithelial neoplasia grade 3 (CIN3). Stage 0 is not included in the FIGO system.

Stage I (T1, N0, M0)

In this stage the cancer has grown into (invaded) the cervix, but it is not growing outside the uterus (T1). The cancer has not spread to nearby lymph nodes (N0) or distant sites (M0).

Stage IA (T1a, N0, M0)

This is the earliest form of stage I. There is a very small amount of cancer, and it can be seen only under a microscope.

The cancer has not spread to nearby lymph nodes (N0) or distant sites (M0).

- **Stage IA1 (T1a1, N0, M0):** The cancer is less than 3 mm (about 1/8-inch) deep and less than 7 mm (about 1/4-inch) wide (T1a1). The cancer has not spread to nearby lymph nodes (N0) or distant sites (M0).
- **Stage IA2 (T1a2, N0, M0):** The cancer is between 3 mm and 5 mm (about 1/5-inch) deep and less than 7 mm (about 1/4-inch) wide (T1a2). The cancer has not spread to nearby lymph nodes (N0) or distant sites (M0).

Stage IB (T1b, N0, M0): This includes stage I cancers that can be seen without a microscope as well as cancers that can only be seen with a microscope if they have spread deeper than 5 mm (about 1/5 inch) into connective tissue of the cervix or are wider than 7 mm (T1b). These cancers have not spread to nearby lymph nodes (N0) or distant sites (M0).

- **Stage IB1 (T1b1, N0, M0):** The cancer can be seen but it is not larger than 4 cm (about 1 3/5 inches) (T1b1). It has not spread to nearby lymph nodes (N0) or distant sites (M0).
- **Stage IB2 (T1b2, N0, M0):** The cancer can be seen and is larger than 4 cm (T1b2). It has not spread to nearby lymph nodes (N0) or distant sites (M0).

Stage II (T2, N0, M0)

In this stage, the cancer has grown beyond the cervix and uterus, but hasn't spread to the walls of the pelvis or the lower part of the vagina. It has not spread to nearby lymph nodes (N0) or distant sites (M0).

Stage IIA (T2a, N0, M0): The cancer has not spread into the tissues next to the cervix (the parametria), but it may have grown into the upper part of the vagina (T2a). It has not spread to nearby lymph nodes (N0) or distant sites (M0).

- **Stage IIA1 (T2a1, N0, M0):** The cancer can be seen but it is not larger than 4 cm (about 1 3/5 inches) (T2a1). It has not spread to nearby lymph nodes (N0) or distant sites (M0).
- **Stage IIA2 (T2a2, N0, M0):** The cancer can be seen and is larger than 4 cm (T2a2). It has not spread to nearby lymph nodes (N0) or distant sites (M0).
- **Stage IIB (T2b, N0, M0):** The cancer has spread into the tissues next to the cervix (the parametria) (T2b). It has not spread to nearby lymph nodes (N0) or distant sites (M0).

Stage III (T3, N0, M0)

The cancer has spread to the lower part of the vagina or the walls of the pelvis, and it may be blocking the ureters (tubes that carry urine from the kidneys to the bladder) (T3). It has not spread to nearby lymph nodes (N0) or distant sites (M0).

- **Stage IIIA (T3a, N0, M0):** The cancer has spread to the lower third of the vagina but not to the walls of the pelvis (T3a). It has not spread to nearby lymph nodes (N0) or distant sites (M0).
 - **Stage IIIB (T3b, N0, M0; OR T1-T3, N1, M0):**
 - The cancer has grown into the walls of the pelvis and/or has blocked one or both ureters causing kidney problems (hydronephrosis) (T3b),
- OR
- The cancer has spread to lymph nodes in the pelvis (N1) but not to distant sites (M0). The tumor can be any size and may have spread to the lower part of the vagina or walls of the pelvis (T1 to T3).

Stage IV

This is the most advanced stage of cervical cancer. The cancer has spread to nearby organs or other parts of the body.

- **Stage IVA (T4, N0, M0):** The cancer has spread to the bladder or rectum, which are organs close to the cervix (T4). It has not spread to nearby lymph nodes (N0) or distant sites (M0).
- **Stage IVB (any T, any N, M1):** The cancer has spread to distant organs beyond the pelvic area, such as the lungs or liver. (M1)

Survival rates for cervical cancer, by stage

Survival rates tell you what percentage of people with the same type and stage of cancer are still alive a certain amount of time (usually 5 years) after they were diagnosed. They can't tell you how long you will live, but they may help give you a better understanding about how likely it is that your treatment will be successful. Some people will want to know the survival rates for their cancer, and some people won't. If you don't want to know, you don't have to.

What is a 5-year survival rate?

Statistics on the outlook for a certain type of cancer are often given as 5-year survival rates. The 5-year survival rate is the percentage of people who live at least 5 years after being diagnosed with cancer. For example, a 5-year survival rate of 70% means that an estimated 70 out of 100 people who have that cancer are still alive 5 years after being diagnosed. Keep in mind, however, that many of these people live much longer than 5 years after diagnosis.

But remember, all survival rates are estimates – your outlook can vary based on a number of factors specific to you.

Cancer survival rates don't tell the whole story

Survival rates are often based on previous outcomes of large numbers of people who had the disease, but they can't predict what will happen in any particular person's case. There are a number of limitations to remember:

- The numbers below are among the most current available. But to get 5-year survival rates, doctors have to look at people who were treated at least 5 years ago. As treatments are improving over time, women who are now being diagnosed with cervical cancer may have a better outlook than these statistics show.
- These statistics are based on when the cancer was first diagnosed. They do not apply to cancers that later come back or spread, for example.
- The outlook for women with cervical cancer varies by the [stage](#) (extent) of the cancer – in general, the survival rates are higher for women with earlier stage cancers. But other factors can also affect a woman's outlook, such as her age and overall health, and how

well the cancer responds to treatment. The outlook for each woman is specific to her circumstances.

Your doctor can tell you how these numbers may apply to you.

Survival rates for cervical cancer

The rates below were published in 2010 in the 7th edition of the AJCC staging manual. They are based on data collected by the National Cancer Data Base from people diagnosed between 2000 and 2002. These are the most recent statistics available for survival by the current staging system.

- The 5-year survival rate for people with stage 0 cervical cancer is about 93%.
- For stage IA cervical cancer, the 5-year survival rate is about 93%. For stage IB cancer, the 5-year survival rate is about 80%.
- For stage IIA cervical cancer, the 5-year survival rate is about 63%. For stage IIB cancer, the 5-year survival rate is about 58%.
- The 5-year survival rate for stage IIIA cervical cancer is about 35%. For stage IIIB cancer, the 5-year survival rate is about 32%.
- Stage IVA cervical cancer has a 5-year survival rate of about 16%, and stage IVB cancer has a 5-year survival rate of about 15%. Still, there are often treatment options available for women with these stages of cancer.

Remember, these survival rates are only estimates – they can't predict what will happen to any individual person. We understand that these statistics can be confusing and may lead you to have more questions. Talk to your doctor to better understand your specific situation.

Treating cervical cancer

If you have been diagnosed with cervical cancer, your cancer care team will discuss your treatment options with you. It's important to think carefully about your choices. You will want to weigh the benefits of each treatment option against the possible risks and side effects.

Although the treatment options depend largely on the stage (extent) of the cancer, other factors may influence your options as well, including your age and general health, your individual circumstances, and your preferences. Cervical cancer can affect your sex life and your ability to have children. These concerns should also be considered as you make treatment decisions. (See [Sex and the Woman With Cancer](#) to learn more about these issues.)

What are cervical cancer treatment options?

Common types of treatments for cervical cancer can include:

- Surgery
- Radiation therapy
- Chemotherapy (chemo)
- Targeted therapy

Some women may need more than one type of treatment.

For the earliest stages of cervical cancer, either surgery or radiation combined with chemo may be used. For later stages, radiation combined with chemo is usually the main treatment. Chemo (by itself) is often used to treat advanced cervical cancer.

Your recovery is the goal of your cancer care team. If a cure is not possible, the goal may be to remove or destroy as much of the cancer as possible to help you live longer and feel better. Sometimes treatment is aimed at relieving symptoms. This is called palliative treatment.

Be sure that you understand all the risks and side effects of the various treatments before making your decision.

What types of doctors treat cervical cancer?

Doctors on your cancer treatment team may include:

- A **gynecologist**: a doctor who treats diseases of the female reproductive system
- A **gynecologic oncologist**: a doctor who treats cancers of the female reproductive system
- A **radiation oncologist**: a doctor who uses radiation to treat cancer

- A **medical oncologist**: a doctor who uses chemotherapy and other medicines to treat cancer

Many other specialists might be part of your treatment team as well, including physician assistants (PAs), nurse practitioners (NPs), nurses, psychologists, social workers, rehabilitation specialists, and other health professionals. See [Health Professionals Associated With Cancer Care](#) for more on this.

Making treatment decisions

It's important to discuss all of your treatment options, including their goals and possible side effects, with your doctors to help make the decision that best fits your needs.

It's also very important to ask questions if there's anything you're not sure about. You can find some good questions to ask in [What should you ask your doctor about cervical cancer?](#)

Getting a second opinion

It is often a good idea to get a second opinion, especially from doctors experienced in treating cervical cancer. A second opinion can give you more information and help you feel more confident about choosing a treatment plan. Some insurance companies require a second opinion before they will agree to pay for certain treatments. Almost all will pay for a second opinion. Still, you might want to check your coverage first, so you'll know if you will have to pay for it.

Thinking about taking part in a clinical trial

Clinical trials are carefully controlled research studies that are done to get a closer look at promising new treatments or procedures. Clinical trials are one way to get state-of-the-art cancer treatment. In some cases, they may be the only way to get access to newer treatments. They are also the best way for doctors to learn better methods to treat cancer. Still, they are not right for everyone.

If you would like to learn more about clinical trials that might be right for you, start by asking your doctor if your clinic or hospital conducts clinical trials. You can also call our clinical trials matching service at 1-800-303-5691 for a list of studies that meet your medical needs, or see [Clinical Trials](#) to learn more.

Considering complementary and alternative methods

You may hear about alternative or complementary methods that your doctor hasn't mentioned to treat your cancer or relieve symptoms. These methods can include vitamins, herbs, and special diets, or other methods such as acupuncture or massage, to name a few.

Complementary methods refer to treatments that are used along with your regular medical care. **Alternative treatments** are used instead of a doctor's medical treatment. Although some of these methods might be helpful in relieving symptoms or helping you feel better, many have not been proven to work to treat the cancer. Some might even be dangerous.

Be sure to talk to your cancer care team about any method you are thinking about using. They can help you learn what is known (or not known) about the method, which can help you make an informed decision. See Complementary and Alternative Medicine to learn more.

Choosing to stop treatment or choosing no treatment at all

For some people, when treatments have been tried and are no longer controlling the cancer, it could be time to weigh the benefits and risks of continuing to try new treatments. Whether or not you continue treatment, there are still things you can do to help maintain or improve your quality of life. Learn more in [If Cancer Treatments Are No Longer Working](#).

Some people, especially if the cancer is advanced, might not want to be treated at all. There are many reasons you might decide not to get cancer treatment, but it's important to talk this through with your doctors before you make that decision. Remember that even if you choose not to treat the cancer, you can still get supportive care to help with pain or other symptoms.

Help getting through cancer treatment

Your cancer care team will be your first source of information and support, but there are other resources for help when you need it. Hospital- or clinic-based support services are an important part of your care. These might include nursing or social work services, financial aid, nutritional advice, rehab, or spiritual help.

The American Cancer Society also has programs and services – including rides to treatment, lodging, support groups, and more – to help you get through treatment. Call our National Cancer Information Center at 1-800-227-2345 and speak with one of our trained specialists on call 24 hours a day, every day.

The treatment information given here is not official policy of the American Cancer Society and is not intended as medical advice to replace the expertise and judgment of your cancer care team. It is intended to help you and your family make informed decisions, together with your doctor. Your doctor may have reasons for suggesting a treatment plan different from these general treatment options. Don't hesitate to ask him or her questions about your treatment options.

What should you ask your doctor about cervical cancer?

It is important for you to have frank, open discussions with your cancer care team. They want to answer all of your questions, so that you can make informed treatment and life decisions. For instance, consider these questions:

When you're told you have cervical cancer

- What type of cervical cancer do I have?
- Has my cancer spread outside of the cervix?
- What is the stage (extent) of my cancer and what does that mean?
- Will I need other tests before we can decide on treatment?
- Do I need to see any other doctors or health professionals?
- If I'm concerned about the costs and insurance coverage for my diagnosis and treatment, who can help me?

When deciding on a treatment plan

- What are my treatment choices?
- What treatment do you recommend and why?
- How much experience do you have treating this type of cancer?
- Should I get a second opinion? How do I do that? Can you recommend someone?
- What would the goal of the treatment be?
- How quickly do we need to decide on treatment?
- What should I do to be ready for treatment?
- How long will treatment last? What will it be like? Where will it be done?
- What risks or side effects are there to the treatments you suggest? Are there things I can do to reduce these side effects?
- How might treatment affect my daily activities?
- What are the chances my cancer will recur (come back) with these treatment plans?

- What will we do if the treatment doesn't work or if the cancer recurs?
- Will I be able to have children after my treatment?
- What are my treatment options if I want to have children in the future?

During treatment

Once treatment begins, you'll need to know what to expect and what to look for. Not all of these questions may apply to you, but asking the ones that do may be helpful.

- How will we know if the treatment is working?
- Is there anything I can do to help manage side effects?
- What symptoms or side effects should I tell you about right away?
- How can I reach you on nights, holidays, or weekends?
- Do I need to change what I eat during treatment?
- Are there any limits on what I can do?
- What kind of exercise should I do, and how often?
- Can you suggest a mental health professional I can see if I start to feel overwhelmed, depressed, or distressed?

After treatment

- Will I need a special diet after treatment?
- Are there any limits on what I can do?
- What other symptoms should I watch for?
- What kind of exercise should I do now?
- What type of follow-up will I need after treatment?
- How often will I need to have follow-up exams and imaging tests?
- Will I need any blood tests?
- How will we know if the cancer has come back? What should I watch for?
- What will my options be if the cancer comes back?

Along with these sample questions, be sure to write down some of your own. For instance, you might want more information about recovery times. Or you might ask if you qualify for a clinical trial .

Keep in mind that doctors aren't the only ones who can give you information. Other health care professionals, such as nurses and social workers, can answer some of your questions. To find out more about speaking with your health care team, see [The Doctor-Patient Relationship](#).

Surgery for cervical cancer

Many women with cervical cancer will have some type of surgery. Surgery can be used to:

- Help diagnose cervical cancer
- Help determine how far the cancer has spread
- Help treat the cancer (especially for earlier-stage cancers)

Several types of 'surgery' can be used to help treat cervical cancer, although some of these destroy cervical tissue (with cold or with a laser) rather than removing it.

Cryosurgery

A very cold metal probe is placed directly on the cervix. This kills the abnormal cells by freezing them. This can be done in a doctor's office or clinic. After cryosurgery, you may have a lot of watery brown discharge for a few weeks.

Laser surgery

A focused laser beam, directed through the vagina, is used to vaporize (burn off) abnormal cells or to remove a small piece of tissue to send to the lab for tests. This can be done in a doctor's office or clinic and is done with local anesthesia (numbing medicine).

Conization

A cone-shaped piece of tissue is removed from the cervix. This is done using a surgical or laser knife (cold knife cone biopsy) or using a thin wire heated by electricity (the loop electrosurgical, LEEP or LEETZ procedure). (See [How are cervical cancers and pre-cancers diagnosed?](#) for more information.) After the procedure, the removed tissue is examined with a microscope. If the margins (outer edges) of the tissue contain cancer (or pre-cancer) cells (called positive margins), some cancer (or pre-cancer) may have been left behind, so further treatment is needed.

Simple (total) hysterectomy

This surgery removes the uterus (both the body of the uterus and the cervix) but not the structures next to the uterus (parametria and uterosacral ligaments). The vagina and pelvic lymph nodes are not removed. The ovaries and fallopian tubes are usually left in place unless there is another reason to remove them.

There are different ways to do a hysterectomy:

- **Abdominal hysterectomy:** The uterus is removed through a surgical incision in the front of the abdomen.
- **Vaginal hysterectomy:** The uterus is removed through the vagina.
- **Laparoscopic hysterectomy:** The uterus is removed using laparoscopy (sometimes called 'keyhole surgery'). A thin tube with a tiny video camera at the end (the laparoscope) is inserted into one or more very small surgical incisions to see inside of the abdomen and pelvis. Small instruments can be controlled through the tube, so the surgeon makes cuts and removes tissue through the tubes without making a large cut in the abdomen.
- **Laparoscopic-assisted vaginal hysterectomy:** The uterus, ovaries, and fallopian tubes are removed through a vaginal incision using the laparoscope, which makes it easier for the doctor.
- **Robotic-assisted surgery:** In this approach, the laparoscopy is done with special tools attached to robotic arms that are controlled by the doctor to help perform precise surgery.

General or epidural (regional) anesthesia is used for all of these operations.

For a laparoscopic or vaginal hysterectomy, the hospital stay is usually 1 to 2 days, followed by a 2- to 3-week recovery period. A hospital stay of 3 to 5 days is common for an abdominal hysterectomy, and complete recovery takes about 4 to 6 weeks.

Possible side effects: Any type of hysterectomy results in infertility (inability to have children). Complications are unusual but could include excessive bleeding, wound infection, or damage to the urinary or intestinal systems.

Hysterectomy does not change a woman's ability to feel sexual pleasure. A woman does not need a uterus or cervix to reach orgasm. The area around the clitoris and the lining of the vagina remain as sensitive as before a hysterectomy. More information about managing the sexual side effects of cervical cancer treatment can be found in *Sex and the Woman with Cancer*.

Radical hysterectomy

For this operation, the surgeon removes the uterus along with the tissues next to the uterus (the parametria and the uterosacral ligaments) and the upper part (about 1 inch) of the vagina next to the cervix. The ovaries and fallopian tubes are not removed unless there is some other medical reason to do so. This surgery is usually done through an abdominal incision. Often, some pelvic lymph nodes are removed as well. (This procedure, known as **lymph node dissection**, is discussed later in this section.)

A radical hysterectomy can also be done using laparoscopy (keyhole surgery). (See the ‘Simple hysterectomy’ section for a description of laparoscopy.) This can result in less pain and a shorter hospital stay compared to surgery using longer incisions.

- **Laparoscopic-assisted radical vaginal hysterectomy** is a surgical approach that combines a radical vaginal hysterectomy with a laparoscopic pelvic lymph node dissection.
- The laparoscope can also make it easier to perform a radical hysterectomy through the abdomen. When lymph nodes are removed in this abdominal procedure, this is called **laparoscopically assisted radical hysterectomy with lymphadenectomy**.
- **Robot-assisted laparoscopic surgery** is also sometimes used for radical hysterectomies. In this approach, the surgeon sits at a control panel to precisely move robotic arms that hold surgical tools. It results in less blood loss during the operation and possibly a shorter hospital stay than other types of laparoscopic surgery, but it’s not clear if long-term outcomes are any better, and its ultimate role in treatment is still being studied.

More tissue is removed in a radical hysterectomy than in a simple one, so the hospital stay can be longer.


Possible side effects: Because the uterus is removed, this surgery results in infertility. Because some of the nerves to the bladder are removed, some women have problems emptying their bladder after this operation and may need a catheter for a time. Complications are unusual but could include excessive bleeding, wound infection, or damage to the urinary and intestinal systems.

Radical hysterectomy does not change a woman's ability to feel sexual pleasure. Although the vagina is shortened, the area around the clitoris and the lining of the vagina is as sensitive as before. A woman does not need a uterus or cervix to reach orgasm. When cancer has caused pain or bleeding with intercourse, the hysterectomy may actually improve a woman's sex life by stopping these symptoms. More information about managing the sexual side effects of cervical cancer treatment can be found in *Sex and the Woman with Cancer*.

Trachelectomy

Another procedure, known as a **radical trachelectomy**, allows women to be treated without losing their ability to have children. The operation is done either through the vagina or the abdomen, and is sometimes done using laparoscopy (keyhole surgery).

This procedure removes the cervix and the upper part of the vagina but not the body of the uterus. The surgeon then places a "purse-string" stitch to act as an artificial opening of the cervix inside the uterine cavity.



The nearby lymph nodes are also removed using laparoscopy, which may require another incision (cut).

After trachelectomy, some women are able to carry a pregnancy to term and deliver a healthy baby by cesarean section, although women who have had this surgery may have a higher risk of miscarriage.

Pelvic exenteration

This is a more extensive operation that may be used to treat recurrent cervical cancer. In this surgery, all of the same organs and tissues are removed as in a radical hysterectomy with pelvic lymph node dissection. (Lymph node dissection is discussed in the next section.) In addition, the bladder, vagina, rectum, and part of the colon may also be removed, depending on where the cancer has spread.

If your bladder is removed, you will need a new way to store and eliminate urine. This usually means using a short segment of intestine to function as a new bladder. The new bladder may be connected to the abdominal wall so that urine is drained periodically when the patient places a catheter into a urostomy (a small opening). Or urine may drain continuously into a small plastic bag attached to the front of the abdomen. For more information see Urostomy: A Guide.

If the rectum and part of the colon are removed, a new way to eliminate solid waste must be created. This is done by attaching the remaining intestine to the abdominal wall so that stool can pass through a colostomy (a small opening) into a small plastic bag worn on the front of the abdomen. (More information about colostomies can be found in *Colostomy: A Guide*.) In some cases, it may be possible to only remove the cancerous part of the colon (next to the cervix) and reconnect the colon ends so that no bags or external appliances are needed.

If the vagina is removed, a new vagina can be surgically created out of skin, intestinal tissue, or muscle and skin (myocutaneous) grafts.

Sexual impact of pelvic exenteration

Recovery from total pelvic exenteration takes a long time. Most women don't begin to feel like themselves again for about 6 months after surgery. Some say it takes a year or two to adjust completely. Nevertheless, these women can lead happy and productive lives. With practice and determination, they can also have sexual desire, pleasure, and orgasms. More information about managing the sexual side effects of cervical cancer treatment can be found in *Sex and the Woman with Cancer*.

Pelvic lymph node dissection

Cancer that starts in the cervix can spread to lymph nodes (pea-sized collections of immune system tissue) in the pelvis. To check for lymph node spread, the surgeon might remove some of these lymph nodes. This procedure is known as a **lymph node dissection** or **lymph node sampling**. It is done at the same time as a hysterectomy or trachelectomy.

Removing lymph nodes can sometimes lead to fluid drainage problems in the leg. This can cause severe swelling in the leg, a condition called **lymphedema**. More information about lymphedema can be found in *Understanding Lymphedema – For Cancers Other Than Breast Cancer*.

Para-aortic lymph node sampling

If cancer is found in any pelvic lymph nodes during surgery, some of the lymph nodes along the aorta (the large artery in the abdomen) may be removed as well. This is called **para-aortic lymph node sampling**. Any tissue removed at surgery will be tested to see if the cancer has spread further than expected. If so, radiation therapy with or without chemotherapy may be recommended.

If surgery is not done, another way to check for lymph node spread is with an imaging test (like MRI or PET/CT). Lymph nodes that are bigger than usual and/or light up on PET scan are more likely to have cancer, so they might need to be biopsied.

Radiation therapy for cervical cancer

Radiation therapy uses high energy x-rays or radioactive particles to kill cancer cells. Radiation therapy may be used for cervical cancer:

- **As a part of the main treatment.** For some stages of cervical cancer, the preferred treatment is radiation alone or surgery followed by radiation. For other stages, radiation and chemo given together (called **concurrent chemoradiation**) is the preferred treatment. The chemo helps the radiation work better.
- **To treat cervical cancer that has spread or that has come back after treatment.** Radiation therapy may be used to treat cancers that have spread to other organs and tissues.

The two types of radiation therapy most often used to treat cervical cancer include:

- External beam radiation
- Brachytherapy

External beam radiation

One way to give radiation is to aim x-rays at the cancer from outside the body. This is called **external beam radiation therapy (EBRT)**. Treatment is much like getting a regular x-ray, but the radiation dose is stronger.

Each radiation treatment lasts only a few minutes, but getting you in the right place in the machine for treatment usually takes longer. The procedure itself is painless.

When radiation is used as the main treatment for cervical cancer, EBRT is usually combined with chemotherapy (called **concurrent chemoradiation**). Often, a low dose of the chemo drug cisplatin is used, but other chemo drugs can be used as well. The radiation treatments are given 5 days a week for a total of 6 to 7 weeks. The chemotherapy is given at scheduled times during the radiation. The schedule is determined by which drug is used.

EBRT can also be used by itself to treat areas of cancer spread or as the main treatment of cervical cancer in patients after surgery or those who can't tolerate chemoradiation.

Possible side effects of EBRT

Side effects of external beam radiation therapy for cervical cancer can include:

- Fatigue (tiredness)
- Upset stomach
- Diarrhea or loose stools

- **Skin changes:** As the radiation passes through the skin to the cancer, it can damage the skin cells. This can cause irritation ranging from mild, short-term redness to peeling. The skin may release fluid, which can lead to infection, so the area exposed to radiation must be carefully cleaned and protected.
- **Radiation cystitis:** Radiation to the pelvis can irritate the bladder (radiation cystitis), causing discomfort and an urge to urinate often.
- **Vaginal pain:** Radiation can make the vulva and vagina more sensitive and sore, and sometimes causes a discharge.
- **Menstrual changes:** Pelvic radiation can affect the ovaries, leading to menstrual changes and even early menopause
- **Low blood counts:** For example, anemia (low levels of red blood cells) can make you feel tired. Leukopenia (low levels of white blood cells increases the risks of serious infection).

When chemotherapy is given with radiation,

the blood counts tend to be lower and fatigue and nausea tend to be worse. These side effects typically improve in the weeks after treatment is stopped.

Other, longer-term side effects are also possible with EBRT. These are described below.

Brachytherapy

Brachytherapy, or internal radiation therapy, puts a source of radiation in or near the cancer. This type of radiation only travels a short distance. The type of brachytherapy used most often to treat cervical cancer is known as **intracavitary brachytherapy**. The radiation source is placed in a device in the vagina (and sometimes in the cervix). This is often used in addition to EBRT as a part of the main treatment for cervical cancer.

There are two types of brachytherapy:

- **Low-dose rate (LDR) brachytherapy** is completed over a few days. During this time, the patient stays in bed in a private room in the hospital with instruments holding the radioactive material in place. While the radiation therapy is being given, the hospital staff will care for you, but also take precautions to lessen their own radiation exposure.
- **High-dose rate (HDR) brachytherapy** is done as an outpatient over several treatments (often at least a week apart). For each high-dose treatment, the radioactive material is inserted for a few minutes and then removed. The advantage of HDR treatment is that you do not have to stay in the hospital or stay still for long periods of time.

To treat cervical cancer in women who have had a hysterectomy, the radioactive material is placed in a tube in the vagina. To treat a woman who still has a uterus, the radioactive

material can be placed in a small metal tube (called a tandem) that goes in the uterus, along with small round metal holders (ovoids) placed near the cervix. This is sometimes called tandem and ovoid treatment. Another option is called tandem and ring. For this, a round holder (like a disc) is placed close to the uterus. The choice of which one to use depends on what type of brachytherapy is planned.

Possible short-term side effects of brachytherapy

Since the radiation only travels a short distance with brachytherapy, the main effects of the radiation are on the cervix and the walls of the vagina. The most common side effect is irritation of the vagina. It may become red and sore, and there may be a discharge. The vulva may become irritated as well.

Brachytherapy can also cause many of the same side effects as EBRT, such as fatigue, diarrhea, nausea, irritation of the bladder, and low blood counts. Often brachytherapy is given right after external beam radiation (before the side effects can go away), so it can be hard to know which type of treatment is causing the side effect.

Long-term side effects of radiation therapy

Vaginal stenosis: Both EBRT and brachytherapy can cause scar tissue to form in the vagina. The scar tissue can make the vagina narrower (called vaginal stenosis), less able to stretch, or even shorter, which can make vaginal sex painful.

A woman can help prevent this problem by stretching the walls of her vagina several times a week, either by having sex or by using a vaginal dilator (a plastic or rubber tube used to stretch out the vagina). For more information, see *Sex and Women With Cancer*.

Vaginal dryness: Vaginal dryness and painful sex can be long-term side effects from radiation (both brachytherapy and EBRT). Estrogens used locally may help with vaginal dryness and changes to the vaginal lining, especially if radiation to the pelvis damaged the ovaries, causing early menopause. These hormones are typically applied into the vagina and absorbed into the genital area, rather than taken by mouth. They come in gel, cream, ring, and tablet forms. For more information, see *Sex and Women With Cancer*.

Weakened bones: Radiation to the pelvis can weaken the bones, leading to fractures. Hip fractures are the most common, and might occur 2 to 4 years after radiation. Bone density tests are recommended to monitor the risk of fracture.

Swelling of the leg(s): If pelvic lymph nodes are treated with radiation, it can lead to fluid drainage problems in the leg. This can cause severe swelling in the leg, a condition called lymphedema. More information about lymphedema can be found in *Understanding Lymphedema – For Cancers Other Than Breast Cancer*.

If you are having side effects from radiation treatment, discuss them with your cancer care team.

It is important to know that smoking increases the side effects from radiation and can make treatment less effective. If you smoke, you should stop.

For more information, see the Radiation Therapy section of our website or A Guide to Radiation Therapy.

Chemotherapy for cervical cancer

Chemotherapy (chemo) uses anti-cancer drugs that are injected into a vein or given by mouth. These drugs enter the bloodstream and can reach all areas of the body, making this treatment useful for killing cancer cells in most parts of the body. Chemo is often given in cycles, with each period of treatment followed by a recovery period. There are a few situations in which chemo may be recommended for cervical cancer.

As a part of the main treatment for cervical cancer

For some stages of cervical cancer, the preferred treatment is radiation and chemo given together (called concurrent chemoradiation). The chemo helps the radiation work better. Options for concurrent chemoradiation include:

- Cisplatin given weekly during radiation. This drug is given into a vein (IV) about 4 hours before the radiation treatment.
- Cisplatin plus 5-fluorouracil (5-FU) given every 4 weeks during radiation.

Sometimes chemo is also given (without radiation) before and/or after chemoradiation.

To treat cervical cancer that has spread or come back after treatment

Chemo may be used to treat cancers that have spread to other organs and tissues. It can also be helpful when cancer comes back after treatment with chemoradiation.

The chemo drugs most often used to treat advanced cervical cancer include:

- Cisplatin
- Carboplatin
- Paclitaxel (Taxol[®])
- Topotecan

- Gemcitabine (Gemzar[®])

Combinations of these drugs are used often.

Some other drugs can be used as well, such as docetaxel (Taxotere[®]), ifosfamide (Ifex[®]), 5-fluorouracil (5-FU), irinotecan (Camptosar[®]), and mitomycin.

The targeted drug bevacizumab (Avastin[®]) may also be added to chemo. This is discussed in the section about targeted therapy.

Side effects of chemotherapy for cervical cancer

Chemo drugs kill cancer cells but also damage some normal cells, which can lead to certain side effects. Side effects depend on the type and dose of the drugs and the length of time you are treated. Common side effects of chemotherapy can include:

- Nausea and vomiting
- Loss of appetite
- Loss of hair
- Mouth sores
- Fatigue (tiredness)
- Diarrhea

Because chemo can damage the blood-producing cells of the bone marrow, the blood cell counts might become low. Low blood cell counts can result in:

- An increased chance of infection (from a shortage of white blood cells)
- Bleeding or bruising after minor cuts or injuries (because of a shortage of blood platelets)
- Shortness of breath (due to low red blood cell counts)

When chemo is given with radiation, the side effects are often more severe. The nausea, fatigue, and problems with low blood counts are often worse. Diarrhea can also be worse if chemo is given at the same time as radiation.

Your health care team will watch for side effects and can give you medicines to help prevent them or treat them to help you feel better. For example, you can be given drugs to help prevent or reduce nausea and vomiting.

Menstrual changes and fertility issues: For younger women who have not had their uterus removed as a part of treatment, changes in menstrual periods are a common side effect of chemo. But even if your periods stop while you are on chemo, you might still be able to get pregnant. Getting pregnant while receiving chemo is not safe, as it could lead to birth defects

and interfere with treatment. This is why it's important that women who are pre-menopausal before treatment and are sexually active discuss with their doctor the options for birth control. Patients who have finished treatment (like chemo) can often go on to have children, but it's important to talk to your doctor about when it is safe to do so.

Premature menopause (not having any more menstrual periods) and infertility (not being able to become pregnant) may occur and may be permanent. Some chemo drugs are more likely to cause this than others. The older a woman is when she gets chemo, the more likely it is that she will become infertile or go through menopause as a result. If this happens, there is an increased risk of bone loss and osteoporosis. Medicines can treat or help prevent problems with bone loss.

Neuropathy: Some drugs used to treat cervical cancer, including paclitaxel and cisplatin, can damage nerves outside of the brain and spinal cord. The injury can sometimes lead to symptoms like numbness, pain, burning or tingling sensations, sensitivity to cold or heat, or weakness, mainly in the hands and feet. This called peripheral neuropathy. In most cases this gets better or even goes away once treatment is stopped, but it might last a long time in some women.

Increased risk of leukemia: Very rarely, certain chemo drugs can damage the genes in bone marrow cells, leading to blood problems and cancers like myelodysplastic syndromes or even acute myeloid leukemia. If this happens, it is usually within 10 years after treatment. In most women, the benefits of chemo in treating the cervical cancer are likely to far outweigh the risk of this serious but rare complication.

Other side effects are also possible. Some of these are more common with certain chemo drugs. Your cancer care team will talk to you about the possible side effects of the specific drugs you are getting. Many side effects go away after treatment is finished, but some can last a long time or even be permanent. It's important to tell your health care team if you have any side effects, as there are often ways to lessen them.

For more information, see the Chemotherapy section of our website, or A Guide to Chemotherapy.

Targeted therapy for cervical cancer

As researchers have learned more about the changes in cancer cells (and nearby cells) that help them grow, they have been able to develop newer drugs that specifically target these changes. These targeted drugs work differently from standard chemotherapy (chemo) drugs and often have different side effects.

For example, for tumors to grow, they must form new blood vessels to keep them nourished. This process is called **angiogenesis**. Some targeted drugs block this new blood vessel growth and are called **angiogenesis inhibitors**.

Bevacizumab (Avastin[®]) is an angiogenesis inhibitor that can be used to treat advanced cervical cancer. It is a monoclonal antibody (a man-made version of a specific immune system protein) that targets vascular endothelial growth factor (VEGF), a protein that helps new blood vessels to form.

This drug is often used with chemo for a time. Then, if the cancer responds, the chemo may be stopped and the bevacizumab given by itself until the cancer starts growing again.

Possible side effects of targeted therapy

The possible side effects of this drug are different from those of chemotherapy drugs. Some of the more common side effects can include:

- High blood pressure
- Feeling tired
- Loss of appetite

Less common but more serious side effects can include:

- Problems with bleeding
- Blood clots
- Wound healing

A rare but serious side effect is the formation of an abnormal connection (called a fistula) between the vagina and part of the colon or intestine.

Targeted Cancer Therapy has more information about the different kinds of drugs considered targeted therapy.

Treatment options for cervical cancer, by stage

The stage of a cervical cancer is the most important factor in choosing treatment. But other factors can also affect your treatment options, including the exact location of the cancer within the cervix, the type of cancer (squamous cell or adenocarcinoma), your age and overall health, and whether you want to have children.

Stage 0 (carcinoma in situ)

Although the AJCC staging system classifies carcinoma in situ (CIS) as the earliest form of cervical cancer, doctors often think of it as a pre-cancer. That is because the cancer cells in CIS are only in the surface layer of the cervix; they have not grown into deeper layers of cells.

All cases of CIS can be cured with the right treatment. However, pre-cancerous changes can sometimes recur (come back) in the cervix or vagina, so it's very important for your doctor to watch you closely after treatment. This includes follow-up with regular Pap tests and in some instances with colposcopy.

For information about work-up and treatment of abnormal Pap test results and cervical pre-cancers other than CIS, see Cervical Cancer Prevention and Early Detection.

Treatment options for squamous cell carcinoma in situ include:

- Cryosurgery
- Laser surgery
- Loop electrosurgical excision procedure (LEEP/LEETZ)
- Cold knife conization
- Simple hysterectomy (as the first treatment or if the cancer returns after other treatments)

Treatment options for adenocarcinoma in situ include:

- Hysterectomy
- Cone biopsy (a possible option for women who wish to have children). The cone specimen must have no cancer cells at the edges, and the woman must be closely watched after treatment. Once the woman has finished having children, a hysterectomy is recommended.

Stage IA1

Treatment for this stage depends on whether or not you want to continue to be able to have children (maintain fertility) and whether or not the cancer has grown into blood or lymph vessels (called lymphovascular invasion).

Treatment options for women who want to maintain fertility:

A **cone biopsy** is the preferred procedure for women who want to have children after the cancer is treated.

- If the edges of the cone don't contain cancer cells (called negative margins), the woman can be watched closely without further treatment as long as the cancer doesn't come back.
- If the edges of the cone biopsy have cancer cells (called positive margins), then cancer may have been left behind. This can be treated with a repeat cone biopsy or a radical trachelectomy (removal of the cervix and upper vagina). A radical trachelectomy is preferred if the cancer has grown into blood or lymph vessels.

Treatment options for women who don't want to maintain fertility:

- A simple (total) hysterectomy may be an option if the cancer shows no lymphovascular invasion.
- If the cancer has grown into blood or lymph vessels, you might need a radical hysterectomy along with removal of the pelvic lymph nodes.

Stage IA2

Treatment for this stage depends in part on whether or not you want to continue to be able to have children (maintain fertility).

Treatment options for women who want to maintain fertility:

- Cone biopsy with removal of pelvic lymph nodes (pelvic lymph node dissection)
- Radical trachelectomy with pelvic lymph node dissection

Treatment options for women who don't want to maintain fertility:

- External beam radiation therapy (EBRT) to the pelvis plus brachytherapy
- Radical hysterectomy with removal of pelvic lymph nodes and sampling of the para-aortic lymph nodes

If none of the lymph nodes are found to have cancer, radiation may still be discussed as an option if the tumor is large, if the tumor has grown into blood or lymph vessels, or if the tumor is invading the surrounding connective tissue that supports organs such as the uterus, bladder, vagina (the stroma).

If the cancer has spread to the tissues next to the uterus (called the parametria) or to any lymph nodes, or if the tissue removed has positive margins, radiation (EBRT) with chemotherapy is usually recommended. The doctor may also advise brachytherapy after the combined chemo and radiation are done.

Stages IB and IIA

The main treatment options are surgery, radiation, or radiation given with chemo (concurrent chemoradiation).

Stages IB1 and IIA1

Treatment options for women who want to maintain fertility:

- Radical trachelectomy with pelvic lymph node dissection

Treatment options for women who don't want to maintain fertility:

- Radical hysterectomy with removal of lymph nodes in the pelvis and some lymph nodes from the para-aortic area

If none of the lymph nodes are found to have cancer, radiation may still be discussed as an option if the tumor is large, if the tumor has grown into blood or lymph vessels, or if the tumor is invading the surrounding connective tissue that supports organs such as the uterus, bladder, vagina (the stroma).

If the cancer has spread to the tissues next to the uterus (called the parametria) or to any lymph nodes, or if the tissue removed has positive margins, radiation (EBRT) with chemotherapy is usually recommended. The doctor may also advise brachytherapy after the combined chemo and radiation are done.

- Radiation using both brachytherapy and external beam radiation therapy may be an option if a woman is not healthy enough for surgery or if she decides they do not want surgery
- Chemotherapy (chemo) may be given with the radiation (concurrent chemoradiation).

Stages IB2 and IIA2

Treatment options:

- Chemoradiation: This is usually the standard treatment. The chemo may be cisplatin or cisplatin plus fluorouracil. The radiation therapy includes both external beam radiation and brachytherapy.
- Radical hysterectomy with pelvic lymph node dissection and para-aortic lymph node sampling: If cancer cells are found in the removed lymph nodes, or in the edges of the tissue removed (positive margins), surgery may be followed by radiation therapy, which is often given with chemo (concurrent chemoradiation).

Some doctors recommend radiation given with chemotherapy first followed by a hysterectomy.

Stages IIB, III and IVA

Treatment options:

- Chemoradiation: The chemo may be cisplatin or cisplatin plus fluorouracil. The radiation therapy includes both external beam radiation and brachytherapy.

Stage IVB

At this stage, the cancer has spread out of the pelvis to other areas of the body. Stage IVB cervical cancer is not usually considered curable. Treatment options include radiation therapy and/or chemo to try to slow the growth of the cancer or help relieve symptoms. Most standard chemo regimens include a platinum drug (cisplatin or carboplatin) along with another drug such as paclitaxel (Taxol), gemcitabine (Gemzar), or topotecan. The targeted drug bevacizumab (Avastin) may be added to chemo.

Clinical trials are testing other combinations of chemo drugs, as well as some other experimental treatments.

Recurrent cervical cancer

Cancer that comes back after treatment is called recurrent cancer. Cancer can come back locally (in or near where it first started, such as the cervix, uterus or nearby the pelvic organs), or it can come back in distant areas (such as the lungs or bone).

If the cancer has recurred in the pelvis only, extensive surgery (pelvic exenteration) may be an option for some patients. Sometimes radiation or chemo may be used to slow the growth of the cancer or help relieve symptoms, but they aren't expected to cure the cancer.

If chemo is used, you should understand the goals and limitations of this therapy. Sometimes chemo can improve your quality of life, and other times it might diminish it. You need to discuss this with your doctor.

New treatments that may benefit patients with distant recurrence of cervical cancer are being evaluated in clinical trials. Clinical Trials may help if you are thinking about participating in a clinical trial.

Cervical cancer during pregnancy

A small number of cervical cancers are found in pregnant women. Most of these (70%) are stage I cancers.

The treatment plan during pregnancy is determined by:

- Tumor size
- If nearby lymph nodes have cancer
- How far along the pregnancy is
- The specific type of cervical cancer

If the cancer is at a very early stage, such as carcinoma in situ (Stage 0) or stage IA, most doctors believe it is safe to continue the pregnancy to term and have treatment several weeks

after birth. Surgery options after birth for early-stage cancers include a hysterectomy, radical trachelectomy, or a cone biopsy.

If the cancer is stage IB or higher, then you and your doctor must decide whether to continue the pregnancy. If not, treatment would be radical hysterectomy and/or radiation. Sometimes chemotherapy can be given during the pregnancy (in the second or third trimester) to shrink the tumor.

If you decide to continue the pregnancy, the baby should be delivered by cesarean section as soon as it is able to survive outside the womb. More advanced cancers typically need to be treated immediately.

The treatment information given here is not official policy of the American Cancer Society and is not intended as medical advice to replace the expertise and judgment of your cancer care team. It is intended to help you and your family make informed decisions, together with your doctor. Your doctor may have reasons for suggesting a treatment plan different from these general treatment options. Don't hesitate to ask him or her questions about your treatment options.

Living as a cervical cancer survivor

For some women with cervical cancer, treatment may remove or destroy the cancer. Completing treatment can be both stressful and exciting. You'll be relieved to finish treatment, yet it's hard not to worry about the cancer coming back. This is very common if you've had cancer.

For other women, [the cancer may never go away](#) completely. These women may get regular treatments with chemotherapy, radiation therapy, or other therapies to try to help keep the cancer in check. Learning to live with cancer that does not go away can be difficult and very stressful.

Life after cervical cancer means returning to some familiar things and also making some new choices.

Ask your doctor for a survivorship care plan

Talk with your doctor about developing a survivorship care plan for you. This plan might include:

- A suggested schedule for follow-up exams and tests
- A schedule for other tests you might need in the future, such as early detection (screening) tests for other types of cancer, or tests to look for long-term health effects from your cancer or its treatment
- A list of possible late- or long-term side effects from your treatment, including what to watch for and when you should contact your doctor

- Diet and physical activity suggestions

Typical follow-up schedules after cervical cancer

Even if you have completed treatment, you will probably have follow-up visits with your doctor for many years. It's very important to go to all of your follow-up appointments. During these visits, your doctors will ask if you are having any problems and may do exams and lab tests or imaging tests to look for signs of cancer or treatment side effects.

Some treatment side effects might last a long time or might not even show up until years after you have finished treatment. Your doctor visits are a good time to ask questions and talk about any changes or problems you notice or concerns you have.

To some extent, the frequency of follow up visits and tests will depend on the stage of your cancer and the chance of it coming back.

Doctor visits

Your doctor will probably recommend you have a physical exam every 3 to 6 months for the first couple of years after treatment, then every 6 months or so for the next few years. People who were treated for early-stage cancers may need exams less often.

Most doctors recommend that women treated for cervical cancer keep getting regular Pap tests no matter how they were treated (surgery or radiation). Although cells for a Pap test are normally from the cervix, if you no longer have a cervix (because you had a trachelectomy or hysterectomy), the cells will be taken from the upper part of the vagina.

Imaging tests

Whether or not your doctor recommends imaging tests will depend on the stage of your cancer and other factors. CT scans may be done if you have worrisome symptoms of the cancer coming back.

Survivors of cervical cancer should also follow the American Cancer Society guidelines for the early detection of cancer, such as those for breast, lung, and colorectal cancer.

Keeping health insurance and copies of your medical records

Even after treatment, it's very important to keep health insurance. Tests and doctor visits cost a lot, and even though no one wants to think of their cancer coming back, this could happen.

At some point after your cancer treatment, you might find yourself seeing a new doctor who doesn't know about your medical history. It's important to keep copies of your medical records to give your new doctor the details of your diagnosis and treatment. Learn more in [Keeping Copies of Important Medical Records](#).

Can I lower my risk of cervical cancer progressing or coming back?

If you have (or have had) cervical cancer, you probably want to know if there are things you can do that might lower your risk of the cancer growing or coming back, such as exercising, eating a certain type of diet, or taking nutritional supplements. At this time, it's not yet clear if those things will help.

It is known that smoking is linked to an increased risk of cervical cancer. While it's not clear if smoking can affect cervical cancer growth or recurrence, it is still helpful to stop smoking to decrease your risk of getting another smoking related cancer (see [Can I get another cancer after having cervical cancer?](#)). Not smoking can also help you tolerate chemotherapy and radiation better and decrease further damage to the cells of the cervix or cervical area.

Adopting other healthy behaviors such as [eating well, getting regular physical activity](#), and staying at a healthy weight might help, but no one knows for sure. However, we do know that these types of changes can have positive effects on your health that can extend beyond your risk of cervical cancer or other cancers.

About dietary supplements

So far, no dietary supplements (including vitamins, minerals, and herbal products) have been shown to clearly help lower the risk of cervical cancer progressing or coming back. This doesn't mean that no supplements will help, but it's important to know that none have been proven to do so.

Dietary supplements are not regulated like medicines in the United States – they do not have to be proven effective (or even safe) before being sold, although there are limits on what they're allowed to claim they can do. If you're thinking about taking any type of nutritional supplement, talk to your health care team. They can help you decide which ones you can use safely while avoiding those that might be harmful.

Can I get another cancer after having cervical cancer?

Cancer survivors can be affected by a number of health problems, but often a major concern is facing cancer again. Cancer that comes back after treatment is called a recurrence. But some cancer survivors may develop a new, unrelated cancer later. This is called a second cancer.

Unfortunately, being treated for cervical cancer doesn't mean you can't get another cancer. Women who have had cervical cancer can still get the same types of cancers that other women get. In fact, they might be at higher risk for certain types of cancer, including:

- Cancers of the mouth and throat
- Cancer of the larynx (voice box)
- Anal cancer
- Vulvar cancer
- Vaginal cancer
- Lung cancer
- Cancers of the bladder and ureter
- Stomach cancer
- Colorectal cancer
- Pancreas cancer

Many of these cancers are linked to smoking and/or infection with the human papilloma virus (HPV), which are also strongly linked to cervical cancer.

The increased risks of acute myeloid leukemia (AML) and cancers of the rectum, bladder, and soft tissue seem to be linked to treatment with radiation.

Can I lower my risk of getting a second cancer?

There are steps you can take to lower your risk and stay as healthy as possible. For example, women who have had cervical cancer should do their best to stay away from tobacco products. Smoking might further increase the risk of some of the second cancers that are more common after cervical cancer.

To help maintain good health, cervical cancer survivors should also:

- Get to and stay at a healthy weight
- Keep physically active
- Eat a healthy diet, with an emphasis on plant foods
- Limit alcohol to no more than 1 drink per day

These steps may also lower the risk of some other health problems.

See [Second Cancers in Adults](#) for more information about causes of second cancers.

Getting emotional support

Some amount of feeling depressed, anxious, or worried is normal when cervical cancer is a part of your life. Some women are affected more than others. But everyone can benefit from help and support from other people, whether friends and family, religious groups, support groups, professional counselors, or others. Learn more in [Emotions After Cancer Treatment](#).

What's new in cervical cancer research?

New ways to prevent and treat cancer of the cervix are being researched. Some of the promising new developments include:

Sentinel lymph node biopsy (SLNB)

During surgery for cervical cancer, lymph nodes in the pelvis may be removed to check for cancer spread. Instead of removing many lymph nodes, a technique called **sentinel lymph node biopsy** can be used to target just the few lymph nodes most likely to contain cancer. In this technique a blue dye containing a radioactive tracer is injected into the cancer and allowed to drain into lymph nodes. Then, during surgery, the lymph nodes that contain radiation and the blue dye can be identified and removed. These are the lymph nodes most likely to contain cancer if it has spread. If these lymph nodes don't contain cancer, the other lymph nodes don't need to be removed. Removing fewer lymph nodes may lower the risk of later problems such as lymphedema of the legs.

SLNB is not a standard procedure for cervical cancer at this time. Available studies suggest that SLNB may be helpful in early-stage cervical cancer, but more studies are planned to see if this procedure should routinely become part of the treatment.

Immunotherapy

In cancer, the immune system cannot control the fast growth of tumor cells. Recently, new drugs called **immune checkpoint inhibitors** have been developed that “reset” the immune system. They have been found to be active in treating a number of types of cancer. Their helpfulness in cervical cancer treatment is not yet known, but clinical trials are underway to find out more. See [Cancer Immunotherapy](#) for more information on this type of treatment.

HPV vaccines

Vaccines have been developed to prevent infection with some of the HPV types associated with cervical cancer. Currently available vaccines are intended to produce immunity to HPV types that cause about 90% of cervical cancers. Studies are being done to see how well these vaccines will reduce the risk of cervical cancer.

Some experimental vaccines are also being studied for women with established HPV infections, to help their immune systems destroy the virus and cure the infection before a cancer develops.

Still other vaccines are meant to help women who already have advanced cervical cancer. These vaccines attempt to produce an immune reaction to the parts of the virus (E6 and E7 proteins) that make the cervical cancer cells grow abnormally. It is hoped that this immunity will kill the cancer cells or stop them from growing. One such study in advanced cervical cancer showed tumor shrinkage with a vaccine against the E7 protein.

Targeted therapy

As researchers have learned more about the gene changes in cells that cause cancer, they have been able to develop newer drugs that specifically target these changes. These targeted drugs work differently from standard chemotherapy drugs. They often have different (and less severe) side effects. These drugs may be used alone or with more traditional chemotherapy.

Pazopanib is a type of targeted drug that blocks certain growth factors that help cancer cells grow. It has shown to be helpful in some early studies of patients with advanced cervical cancer. This drug continues to be studied.

Hyperthermia

Hyperthermia is a treatment that raises the temperature in the area where the tumor is located, most often by using radiofrequency rods placed around the patient. Some research indicates that adding hyperthermia to radiation may help keep the cancer from coming back and help patients live longer.

Additional resources for cervical cancer

More information from your American Cancer Society

We have a lot more information that you might find helpful. Explore www.cancer.org or call our National Cancer Information Center toll-free number, 1-800-227-2345. We're here to help you any time, day or night.

Other national organizations and websites*

Along with the American Cancer Society, other sources of information and support include:

National Cancer Institute (NCI)

Toll-free number: 1-800-422-6237 (1-800-4-CANCER)

Website: www.cancer.gov

Their “Cancer Information Service” offers a wide variety of free, accurate, up-to-date information about cancer to patients, their families, and the general public; also can help people find clinical trials in their area.

National Women's Health Information Center (NWHIC)

Toll-free number: 1-800-994-9662 (1-800-994-WOMAN)

TTY: 1-888-220-5446

Website: www.womenshealth.gov

Offers a lot of information on women's health issues – including cancers in women

National Cervical Cancer Coalition

Toll-free number: 1-800- 685-5531

Website: www.nccc-online.org

Provides referrals to uninsured, underserved women; information about cervical cancer and its treatment; and phone and email support services

Planned Parenthood Federation of America

Toll-free number: 1-800-230-7526

Website: www.plannedparenthood.org

Offers many women’s health services including cervical cancer screening with referrals for specialty care. Services available for the uninsured.

Foundation for Women’s Cancer (formerly the Gynecologic Cancer Foundation)

Toll-free number: 1-800-444-4441

Telephone number: 1-312-578-1439

Website: www.foundationforwomenscancer.org/

Has a directory of specially trained gynecologic oncologists practicing in the US; free information; and an online "survivor section" featuring articles on personal issues such as fertility, sexuality and quality of life aimed at creating an online community for women with cancer.

National Coalition for Cancer Survivorship (NCCS)

Toll-free number: 1-877-622-7937 (1-877-NCCS-YES)

Website: www.canceradvocacy.org

Has publications on many topics, including employment and health insurance, as it relates to cancer. Materials are also offered in Spanish. Also offers the Cancer Survival Toolbox – a free program that teaches skills to help people with cancer meet the challenges of their illness.

**Inclusion on this list does not imply endorsement by the American Cancer Society.*

No matter who you are, we can help. Contact us anytime, day or night, for information and support. Call us at **1-800-227-2345** or visit www.cancer.org.

References: Cervical cancer detailed guide

Adam E, Kaufman RH, Adler-Storthz K, et al. A prospective study of association of herpes simplex virus and human papillomavirus infection with cervical neoplasia in women exposed to diethylstilbestrol in utero. *Int J Cancer*. 1985;35:19-26.

American Cancer Society. *Cancer Facts and Figures 2016*. Atlanta, Ga: American Cancer Society; 2016.

American Joint Committee on Cancer. Cervix Uteri. In: *AJCC Cancer Staging Manual*. 7th ed. New York, NY: Springer; 2010:395-402.

Anderson TA, Schick V, Herbenick D, Dodge B, Fortenberry JD. A study of human papillomavirus on vaginally inserted sex toys, before and after cleaning, among women who have sex with women and men. *Sex Transm Infect*. 2014 Apr 16. [Epub ahead of print]

Arbyn M, Kyrgiou M, Simoens C, et al. Perinatal mortality and other severe adverse pregnancy outcomes associated with treatment of cervical intraepithelial neoplasia: meta-analysis. *BMJ*. 2008;337: a1284.

Ault KA. Future II study group. Effect of prophylactic human papillomavirus L1 virus-like-particle vaccine on risk of cervical intraepithelial neoplasia grade 2, grade 3, and adenocarcinoma in situ: A combined analysis of four randomised clinical trials. *Lancet*. 2007;369:1861-1868.

Bentivegna E, Maulard A, Pautier P, et al. Fertility results and pregnancy outcomes after conservative treatment of cervical cancer: A systematic review of the literature. *Fertil Steril*. 2016;106:1195-1211.

Castellsagué X, Bosch FX, Muñoz N, et al. Male circumcision, penile human papilloma virus infection, and cervical cancer in female partners. *N Engl J Med*. 2002;346:1105-1112.

Castellsagué X, Díaz M, Vaccarella S, et al. Intrauterine device use, cervical infection with human papillomavirus, and risk of cervical cancer: A pooled analysis of 26 epidemiological studies. *Lancet Oncol*. 2011;12:1023-1031.

Chaturvedi AK, Engels EA, Gilbert ES, et al. Second cancers among 104760 survivors of cervical cancer: Evaluation of long-term risk. *J Natl Cancer Inst.* 2007; 99:1634-1643.

Chaturvedi AK, Kleinerman RA, Hildesheim A, et al. Second cancers after squamous cell carcinoma and adenocarcinoma of the cervix. *J Clin Oncol.* 2009;27:967-973.

Cuzick J, Myers O, Hunt WC, et al. A population-based evaluation of cervical screening in the United States: 2008-2011. *Cancer Epidemiol Biomarkers Prev.* 2014;23:765-73.

Del Priore G, Gudipudi DK, Montemarano N, et al. Oral diindolylmethane (DIM): pilot evaluation of a nonsurgical treatment for cervical dysplasia. *Gynecol Oncol.* 2010;116:464-467.

Klopp AH, Eifel PJ, Berek JS, Konstantinopoulos PA. Chapter 72: Cancer of the cervix, vagina, and vulva. In: DeVita VT, Hellman S, Rosenberg SA, eds. *Cancer: Principles and Practice of Oncology* 10th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2015.

Ghosh C, Baker JA, Moysich KB, et al. Dietary intakes of selected nutrients and food groups and risk of cervical cancer. *Nutr Cancer.* 2008;60:331-341.

Gray RH, Serwadda D, Kong X, et al. Male circumcision decreases acquisition and increases clearance of high-risk human papillomavirus in HIV-negative men: A randomized trial in Rakai, Uganda. *J Infect Dis.* 2010;201:1455-1462.

Grimm C, Polterauer S, Natter C, et al. Treatment of cervical intraepithelial neoplasia with topical imiquimod: a randomized controlled trial. *Obstet Gynecol.* 2012;120:152-159.

Hatch EE, Herbst AL, Hoover RN, et al. Incidence of squamous neoplasia of the cervix and vagina in women exposed prenatally to diethylstilbestrol (United States). *Cancer Causes Control.* 2001;12:837-845.

Hernandez BY, Wilkens LR, Zhu X, et al. Transmission of human papillomavirus in heterosexual couples. *Emerg Infect Dis.* 2008;14:888-894.

Howlander N, Noone AM, Krapcho M, et al (eds). SEER Cancer Statistics Review, 1975-2013, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2013/, based on November 2015 SEER data submission, posted to the SEER web site, April 2016.

International Collaboration of Epidemiological Studies of Cervical Cancer. Cervical cancer and reproductive factors: Collaborative reanalysis of individual data on 16,563 women with cervical carcinoma and 33,542 women without cervical carcinoma from 25 epidemiological studies. *Int J Cancer.* 2006; 119:1108-1124.

International Collaboration of Epidemiological Studies of Cervical Cancer. Appleby P, Beral V, Berrington de González A, et al. Cervical cancer and hormonal contraceptives: Collaborative reanalysis of individual data for 16,573 women with cervical cancer and 35,509 women without cervical cancer from 24 epidemiological studies. *Lancet.* 2007;370:1609-1621.

Jhingran A, Russel AH, Seiden MV, et al. Cancers of the cervix, vagina and vulva. In: Abeloff MD, Armitage JO, Lichter AS, et al. *Clinical Oncology*. 5th ed. Philadelphia, Pa; Elsevier; 2008: 1534-1574.

Kataja V, Syrjänen S, Yliskoski M, et al. Risk factors associated with cervical human papillomavirus infections: a case-control study. *Am J Epidemiol*. 1993;138:735-745.

Kleinerman RA, Kosary C, Hildesheim A. New Malignancies Following Cancer of the Cervix Uteri, Vagina, and Vulva. In: Curtis RE, Freedman DM, Ron E, Ries LAG, Hacker DG, Edwards BK, Tucker MA, Fraumeni JF Jr. (eds). *New Malignancies Among Cancer Survivors: SEER Cancer Registries, 1973-2000*. National Cancer Institute. NIH Publ. No. 05-5302. Bethesda, MD, 2006. Accessed at http://seer.cancer.gov/archive/publications/mpmono/MPMonograph_complete.pdf on November 8, 2016.

Kosary CL. Cancer of the uterine cervix. In: Ries LAG, Young JL, Keel GE, Eisner MP, Lin YD, Horner M-J (editors). *SEER Survival Monograph: Cancer Survival Among Adults: U.S. SEER Program, 1988-2001, Patient and Tumor Characteristics*. National Cancer Institute, SEER Program, NIH Pub. No. 07-6215, Bethesda, MD, 2007.

Kreimer AR, González P, Katki HA, et al. Efficacy of a bivalent HPV 16/18 vaccine against anal HPV 16/18 infection among young women: A nested analysis within the Costa Rica Vaccine Trial. *Lancet Oncol*. 2011;12:862-870.

Kushi LH, Doyle C, McCullough M, Rock CL, et al. American Cancer Society Guidelines on nutrition and physical activity for cancer prevention: Reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J Clin*. 2012;62:30-67.

Kyrgiou M, Koliopoulos G, Martin-Hirsch P, et al. Obstetric outcomes after conservative treatment for intraepithelial or early invasive cervical lesions: Systematic review and meta-analysis. *Lancet*. 2006;367:489-498.

Lacey JV Jr, Swanson CA, Brinton LA, et al. Obesity as a potential risk factor for adenocarcinomas and squamous cell carcinomas of the uterine cervix. *Cancer*. 2003;98:814-821.

Lécuru F, Mathevet P, Querleu D, et al. Bilateral negative sentinel nodes accurately predict absence of lymph node metastasis in early cervical cancer: results of the SENTICOL study. *J Clin Oncol*. 2011;29:1686-1691.

Lu B, Wu Y, Nielson CM, et al. Factors associated with acquisition and clearance of human papillomavirus infection in a cohort of US men: A prospective study. *J Infect Dis*. 2009;199:362-371.

Lutgens L, van der Zee J, Pijls-Johannesma M, et al. Combined use of hyperthermia and radiation therapy for treating locally advanced cervix carcinoma. *Cochrane Database Syst Rev*. 2010;3:CD006377

- Marrazzo JM, Koutsky LA, Stine KL, et al. Genital human papillomavirus infection in women who have sex with women. *J Infect Dis.* 1998;178:1604-1609.
- Massad LS, Einstein MH, Huh WK, et al. 2012 Updated Consensus Guidelines for the Management of Abnormal Cervical Cancer Screening Tests and Cancer Precursors. *J Low Genit Tract Dis.* 2013;17:S1-S27.
- Monk BJ, Sill MW, Burger RA, et al. Phase II trial of bevacizumab in the treatment of persistent or recurrent squamous cell carcinoma of the cervix: A gynecologic oncology group study. *J Clin Oncol.* 2009;27:1069-1074.
- Monk BJ, Mas Lopez L, Zarba JJ, et al. Phase II, open-label study of pazopanib or lapatinib monotherapy compared with pazopanib plus lapatinib combination therapy in patients with advanced and recurrent cervical cancer. *J Clin Oncol.* 2010;28:3562-3569.
- National Institutes of Health. Medline Plus. Female Condoms. 04/05/2016. Accessed at www.nlm.nih.gov/medlineplus/ency/article/004002.htm on Dec 04, 2016.
- NCCN Practice Guidelines in Oncology. Cervical Cancer Version 1.2017. Accessed at www.nccn.org on October 10, 2016.
- Nielson CM, Harris RB, Flores R, et al. Multiple-type human papillomavirus infection in male anogenital sites: prevalence and associated factors. *Cancer Epidemiol Biomarkers Prev.* 2009;18(4):1077-1083. Epub 2009 Mar 24.
- Paavonen J, Naud P, Salmerón J, et al. Efficacy of human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine against cervical infection and precancer caused by oncogenic HPV types (PATRICIA): final analysis of a double-blind, randomised study in young women. *Lancet.* 2009;374:301-314.
- Park DA, Yun JE, Kim SW, Lee SH. Surgical and clinical safety and effectiveness of robot-assisted laparoscopic hysterectomy compared to conventional laparoscopy and laparotomy for cervical cancer: A systematic review and meta-analysis. *Eur J Surg Oncol.* 2016;S0748-7983(16)30687-4.
- PDQ database. Cervical cancer. Bethesda, Md: National Cancer Institute; 02/04/2016. Accessed at www.cancer.gov/cancertopics/pdq/treatment/cervical/healthprofessional on Dec 4, 2016.
- Plummer M, Herrero R, Franceschi S, et al; IARC Multi-centre Cervical Cancer Study Group. Smoking and cervical cancer: Pooled analysis of the IARC multi-centric case-control study. *Cancer Causes Control.* 2003;14:805-814.
- Ronco G, Cuzick J, Pierotti P, et al. Accuracy of liquid based versus conventional cytology: overall results of new technologies for cervical cancer screening: Randomised controlled trial. *BMJ.* 2007;335(7609):28. Epub 2007 May 21.

Rose PG, Ali S, Watkins E, et al. Long-term follow-up of a randomized trial comparing concurrent single agent cisplatin, cisplatin-based combination chemotherapy, or hydroxyurea during pelvic irradiation for locally advanced cervical cancer: A Gynecologic Oncology Group Study. *J Clin Oncol*. 2007;25:2804-2810.

Sadalla JC, Moreira de Andrade J, et al. Cervical cancer: What's new? *Rev Assoc Med Bras* 2015; 61(6):536-542.

Sagae S, Monk BJ, et al. Advances and concepts in cervical cancer trials: A road map for the future. *Int J Gynecol Cancer*. 2016;26:199-207.

Saslow D, Solomon D, Lawson H, et al. American Cancer Society, American Society for Colposcopy and Cervical Pathology, and American Society for Clinical Pathology screening guidelines for the prevention and early detection of cervical cancer. *CA Cancer J Clin*. 2012;62:147-172.

Saslow D, Castle PE, Cox JT, et al. American Cancer Society guideline for human papillomavirus (HPV) vaccine use to prevent cervical cancer and its precursors. *CA Cancer J Clin*. 2007; 57:7-28.

Schiffman M, Castle PE, Jeronimo J, Rodriguez AC, Wacholder S. Human papillomavirus and cervical cancer. *Lancet*. 2007;370:890-907.

Schover LR. *Sexuality and Fertility After Cancer*. New York: Wiley; 1997.

Sellers JW, Mahony JB, Kaczorowski J, et al. Prevalence and predictors of human papillomavirus infection in women in Ontario, Canada. Survey of HPV in Ontario Women (SHOW) Group. *CMAJ*. 2000;163:503-508.

Shepherd JH, Spencer C, Herod J, Ind TEJ. Radical vaginal trachelectomy as a fertility-sparing procedure in women with early-stage cervical cancer--cumulative pregnancy rate in a series of 123 women. *BJOG*. 2006; 113:719-723.

Skrzypczyk-Ostaszewicz A, Rubach, M. Gynaecological cancers coexisting with pregnancy – a literature review. *Contemp Oncol (Pozn)*. 2016;20:193–198.

Tewari KS, Sill MW, Long HJ 3rd, et al. Improved survival with bevacizumab in advanced cervical cancer. *N Engl J Med*. 2014;370:734-743.

Tobian AA, Serwadda D, Quinn TC, et al. Male circumcision for the prevention of HSV-2 and HPV infections and syphilis. *N Engl J Med*. 2009;360:1298-1309.

Tokudome S, Suzuki S, Ichikawa H, et al. Condom use promotes regression of cervical intraepithelial neoplasia and clearance of human papillomavirus: a randomized clinical trial. *Int J Cancer*. 2004;112:164.

Troisi R, Hatch EE, Titus-Ernstoff L, et al. Cancer risk in women prenatally exposed to diethylstilbestrol. *Int J Cancer*. 2007;121:356-360.

Waggoner SE. Cervical cancer. *Lancet*. 2003; 361:2217-2225.

Wheeler CM, Castellsagué X, Garland SM, et al. Cross-protective efficacy of HPV-16/18 AS04-adjuvanted vaccine against cervical infection and precancer caused by non-vaccine oncogenic HPV types: 4-year end-of-study analysis of the randomised, double-blind PATRICIA trial. *Lancet Oncol*. 2012;13:100-110.

Willows K, Lennox G, Covens A. Fertility-sparing management in cervical cancer: balancing oncologic outcomes with reproductive success. *Gynecol Oncol Res Pract*. 2016; Oct 21; 3:9. eCollection.

Winer RL, Hughes JP, Feng Q, et al. Condom use and the risk of genital human papillomavirus infection in young women. *N Engl J Med*. 2006;354:2645-2654.

Winer RL, Lee SK, Hughes JP, et al. Genital human papillomavirus infection: incidence and risk factors in a cohort of female university students. *Am J Epidemiol*. 2003;157(3):218-226. Erratum in: *Am J Epidemiol*. 2003;157(9):858.

Wright TC Jr, Massad LS, Dunton CJ, et al; 2006 American Society for Colposcopy and Cervical Pathology-sponsored Consensus Conference. 2006 consensus guidelines for the management of women with abnormal cervical cancer screening tests. *Am J Obstet Gynecol*. 2007;197:346-355.

Zamarin D, Jazaeri AA. Leveraging immunotherapy for the treatment of gynecologic cancers in the era of precision medicine. *Gynecol Oncol*. 2016;141:86-94.

Last Medical Review: 9/19/2014

Last Revised: 1/29/2016

2014 Copyright American Cancer Society

For additional assistance please contact your American Cancer Society
1-800-227-2345 or www.cancer.org